

AMS-02 Antiprotons Reloaded

Rolf Kappl

Bethe Center for Theoretical Physics & Physikalisches Institut der Universität
Bonn

based on RK, Annika Reinert, Martin Wolfgang Winkler [arXiv:1506.04145](https://arxiv.org/abs/1506.04145)

Anticipating Discoveries: LHC14 and Beyond,
July 14th 2015

Outline

Motivation

Cosmic Ray Propagation


Antiprotons Reloaded – Dark Matter Under Siege

Conclusions

AMS-02 – A Precision Experiment For Cosmic Rays




Primaries vs. Secondaries – Dark Matter?




- Primaries $\xrightarrow{\text{Spallation}}$ Secondaries
- Dark matter annihilation

Primaries vs. Secondaries – Dark Matter?




- Primaries $\xrightarrow{\text{Spallation}}$ Secondaries
- Dark matter annihilation

Primaries vs. Secondaries – Dark Matter?




- Primaries $\xrightarrow{\text{Spallation}}$ Secondaries
- Dark matter annihilation

Primaries vs. Secondaries – Dark Matter?



- Primaries $\xrightarrow{\text{Spallation}}$ Secondaries
- Dark matter annihilation


Transport Of Cosmic Rays – Fokker-Planck



- Two-zone diffusion model

$$\nabla(-K \nabla N_i + \mathbf{V}_c N_i) + \partial_E(b_{\text{tot}} N_i - K_{EE} \partial_E N_i) + \Gamma_{\text{ann}} N_i = q_i$$


Transport Of Cosmic Rays – Fokker-Planck



- Two-zone diffusion model

$$\nabla \cdot (-K \nabla N_i + \mathbf{v}_c N_i) + \partial_E (b_{\text{tot}} N_i - K_{EE} \partial_E N_i) + \Gamma_{\text{ann}} N_i = q_i$$


Transport Of Cosmic Rays – Fokker-Planck



- Two-zone diffusion model

$$\nabla(-K \nabla N_i + \mathbf{v}_c N_i) + \partial_E(b_{\text{tot}} N_i - K_{EE} \partial_E N_i) + \Gamma_{\text{ann}} N_i = q_i$$


Transport Of Cosmic Rays – Fokker-Planck



- Two-zone diffusion model

$$\nabla(-K \nabla N_i + \mathbf{V}_c N_i) + \partial_E(b_{\text{tot}} N_i - K_{EE} \partial_E N_i) + \Gamma_{\text{ann}} N_i = q_i$$

Transport Of Cosmic Rays – Fokker-Planck



- Two-zone diffusion model

$$\nabla(-K \nabla N_i + \mathbf{v}_c N_i) + \partial_E(b_{\text{tot}} N_i - K_{EE} \partial_E N_i) + \Gamma_{\text{ann}} N_i = q_i$$

Bethe's Legacy – Boron As A Tracer



Bethe Center for Theoretical Physics

- CNO cycle shows that boron is not produced in stars
- Boron is produced by spallation of interstellar matter
- Injected boron $q_B \xrightarrow{\text{Propagation}}$ Measured boron at AMS-02
- Task:

$$q_B = 4\pi \sum_{i \in \{C, N, O, Ne, Mg, Si\}} \int dT' \frac{d\sigma}{dT} n_{ISM} \Phi_i(T')$$

MARCH 1, 1939

PHYSICAL REVIEW

VOLUME 55

Energy Production in Stars*

H. A. BETHE

Cornell University, Ithaca, New York

(Received September 7, 1938)

It is shown that the most important source of energy in ordinary stars is the reactions of carbon and nitrogen with protons. These reactions form a cycle in which the original nucleus is reproduced, viz. $C^{12}+H=N^{13}$, $N^{13}=C^{13}+e^+$, $C^{13}+H=N^{14}$, $N^{14}+H=O^{16}$, $O^{16}=N^{15}+e^+$, $N^{15}+H=C^{12}+He^4$. Thus carbon and nitrogen merely serve as catalysts for the combination of four protons (and two electrons) into an α -particle (§7).

The carbon-nitrogen reactions are unique in their vertical character (§8). For all nuclei lighter than carbon.

integration of the Eddington equations gives 19. For the brilliant star Y Cygni the corresponding figures are 30 and 32. This good agreement holds for all bright stars of the main sequence, but, of course, not for giants.

For fainter stars, with lower central temperatures, the reaction $H+H=D+e^+$ and the reactions following it, are believed to be mainly responsible for the energy production. (§10)

It is shown further (§5-6) that no elements heavier than He⁴ can be built up in ordinary stars. This is due to the fact.

Bethe's Legacy – Boron As A Tracer



Bethe Center for Theoretical Physics

- CNO cycle shows that boron is not produced in stars
- Boron is produced by spallation of interstellar matter
- Injected boron $q_B \xrightarrow{\text{Propagation}}$ Measured boron at AMS-02
- Task:

$$q_B = 4\pi \sum_{i \in \{C, N, O, Ne, Mg, Si\}} \int dT' \frac{d\sigma}{dT} n_{ISM} \Phi_i(T')$$

MARCH 1, 1939

PHYSICAL REVIEW

VOLUME 55

Energy Production in Stars*

H. A. BETHE

Cornell University, Ithaca, New York

(Received September 7, 1938)

It is shown that the most important source of energy in ordinary stars is the reactions of carbon and nitrogen with protons. These reactions form a cycle in which the original nucleus is reproduced, viz. $C^{12}+H=N^{13}$, $N^{13}=C^{13}+e^+$, $C^{13}+H=N^{14}$, $N^{14}+H=O^{16}$, $O^{16}=N^{16}+e^+$, $N^{16}+H=C^{12}+He^4$. Thus carbon and nitrogen merely serve as catalysts for the combination of four protons (and two electrons) into an α -particle (§7).


The carbon-nitrogen reactions are unique in their vertical character (§8). For all nuclei lighter than carbon.

integration of the Eddington equations gives 19. For the brilliant star Y Cygni the corresponding figures are 30 and 32. This good agreement holds for all bright stars of the main sequence, but, of course, not for giants.

For fainter stars, with lower central temperatures, the reaction $H+H=D+e^+$ and the reactions following it, are believed to be mainly responsible for the energy production. (§10)

It is shown further (§5-6) that no elements heavier than He⁴ can be built up in ordinary stars. This is due to the fact.

Cosmic Ray Primaries




- We considered C, N, O, Ne, Mg, Si as the main contributions for boron production
- Fits to measured fluxes

$$\Phi_i(T) = A \left(\frac{T}{T+b} \right)^c T^{-\gamma}$$

- Yet no AMS-02 data
- Solar modulation

Cosmic Ray Primaries




- We considered C, N, O, Ne, Mg, Si as the main contributions for boron production
- Fits to measured fluxes

$$\Phi_i(T) = A \left(\frac{T}{T + b} \right)^c T^{-\gamma}$$

- Yet no AMS-02 data
- Solar modulation

Interlude – Cosmic Rays In The Heliosphere



- Force field approximation for solar modulation

$$\Phi^{\text{TOA}}(T - Z\phi) = \frac{(T - Z\phi)^2 + 2m(T - Z\phi)}{T^2 + 2mT} \Phi^{\text{IS}}(T)$$


- Force field parameter ϕ takes care of the solar activity

Back To The Injection – Cross Sections Reloaded

- Source term:

$$q_B = 4\pi \sum_{i \in \{C, N, O, Ne, Mg, Si\}} \int dT' \left[\frac{d\sigma}{dT} \right] n_{ISM} \Phi_i(T')$$

- Parametrization from [Webber et al., AJSS 144 \(2003\) 153](#)
- Large uncertainties, poor experimental data




Nuclear Physics Meets Astrophysics

- Source term q_B for boron determined
- Fokker-Planck equation solved with semi-analytical approach in two dimensions [Maurin et al., Astrophys.J. 555 \(2001\) 585-596](#)
- With the fit to the carbon flux \Rightarrow B/C ratio computed (Secondary/Primary ratio)
- Scanned over a sample of random propagation parameters and kept only configurations which passed χ^2 test


Propagation parameters fixed with highest precision!

B/C Before AMS-02




- Best data before AMS-02 from the PAMELA experiment
[Adriani et al., Astrophys. J. 791 \(2014\), 93](#)
- Clear trend but still large uncertainties

B/C In The AMS-02 Precision Era




- Low experimental errors and data up to high rigidities
- Still preliminary data \Rightarrow Data quality will further increase

\bar{p}/p Ratio With Old Propagation Parameters




- \bar{p}/p ratio computed with outdated (unknown?) propagation parameters

\bar{p}/p Ratio With New Propagation Parameters





- \bar{p}/p ratio computed with new cross sections ([Martin's talk](#))
- New propagation parameters derived from boron \Rightarrow Good agreement for antiprotons
- No need for antiprotons from dark matter annihilation \Rightarrow Good probe to constrain dark matter models

Outlook – Positrons, Better Cross Section Data, . . .





- Data on positrons can be used to further constrain propagation parameters

Outlook – Positrons, Better Cross Section Data, . . .



- New cross section measurements would dramatically decrease the uncertainty

Outlook – Positrons, Better Cross Section Data, . . .



- New cross section measurements would dramatically decrease the uncertainty

Conclusions

- With AMS-02 the precision era for indirect dark matter detection started
- We estimated the antiproton background with very high precision
- The new antiproton data together with our background calculation will be a perfect probe for dark matter models

Thank you for your attention!