

Heavy Flavors and Exotic Hadron Spectroscopy

Joan Soto

The Present and Future of Heavy Flavour and Exotic Hadron Spectroscopy, MIAPbP,
30/05/23



UNIVERSITAT DE
BARCELONA

Today: Ruben Oncala, JS, Phys. Rev. D **96**, 014004 (2017)
JS, Jaume Tarrús Castellà, Phys. Rev. D **102**, 014012 (2020)
JS, Sandra Tomàs Valls, e-Print:2302.01765

Thu: JS, Jaume Tarrús Castellà, Phys. Rev. D **102**, 014013 (2020) ; D
104 (2021) 074027

Heavy Flavors

- Heavy quarks: $Q = c, b, t, m_Q \gg \Lambda_{QCD}$
- Heavy hadrons: hadrons containing at least a heavy quark: $Q = b, c$
- In the hadron rest frame the heavy quark moves slowly \implies use a non-relativistic approximation
- A universal way to encode it together with relativistic correction is using Effective Field Theories
- NRQCD/HQET are the suitable ones
- They imply heavy quark spin symmetry at leading order.

NRQCD

W.E. Caswell and G.P. Lepage, Phys. Lett. **167B**, 437 (1986)

G. T. Bodwin, E. Braaten and G. P. Lepage, Phys. Rev. D **51** (1995) 1125

$$m_Q \gg m_Q v, \quad m_Q v^2, \quad \Lambda_{QCD}$$

$$\mathcal{L}_\psi = \psi^\dagger \left\{ iD_0 + \frac{1}{2m_Q} \mathbf{D}^2 + \frac{1}{8m_Q^3} \mathbf{D}^4 + \frac{c_F}{2m_Q} \boldsymbol{\sigma} \cdot g\mathbf{B} + \frac{c_D}{8m_Q^2} (\mathbf{D} \cdot g\mathbf{E} - g\mathbf{E} \cdot \mathbf{D}) + i \frac{c_S}{8m_Q^2} \boldsymbol{\sigma} \cdot (\mathbf{D} \times g\mathbf{E} - g\mathbf{E} \times \mathbf{D}) \right\} \psi$$

c_F , c_D and c_S are short distance matching coefficients calculable from QCD in powers of α_s . They depend on m_Q and μ (factorization scale) but not on the lower energy scales.

Exotic Hadrons

- Hadrons beyond mesons $q\bar{q}$ and baryons qqq
- QCD: any color singlet state made out of quarks and gluons may become a hadron
- I will restrict myself to discuss hadrons containing two heavy quarks
- The starting point can then be NRQCD

Hadrons with two heavy quarks

$$Q = b, c \quad , \quad q = u, d, s$$

- QQ + light quarks and gluons

- ▶ Double Heavy Baryons: QQq
- ▶ Tetraquarks: $QQ\bar{q}\bar{q}$
- ▶ Pentaquarks: $QQqq\bar{q}$
- ▶ Hybrids: $QQqg$
- ▶ ...

- $Q\bar{Q}$ + light quarks and gluons


- ▶ Heavy Quarkonium: $Q\bar{Q}$
- ▶ Hybrids: $Q\bar{Q}g$
- ▶ Tetraquarks: $Q\bar{Q}q\bar{q}$
- ▶ Pentaquarks: $Q\bar{Q}qqq$
- ▶ ...

Heavy Quarkonium

$Q\bar{Q}$ bound state , $m_Q \gg \Lambda_{QCD}$, $\alpha_s(m_Q) \ll 1$


- Heavy quarks move slowly $v \ll 1$
- Non-relativistic system \rightarrow multiscale problem
 - ▶ $m_Q \gg m_Q v$ (Relative momentum)
 - ▶ $m_Q v \gg m_Q v^2$ (Binding energy)
 - ▶ $m_Q \gg \Lambda_{QCD}$
- EFTs are useful (N. Brambilla, A. Pineda, JS and A. Vairo, Rev. Mod. Phys. 77, 1423 (2005))
 - ▶ NRQCD: $m_Q \gg m_Q v, m_Q v^2, \Lambda_{QCD}$ (W.E. Caswell and G.P. Lepage, Phys. Lett. **167B**, 437 (1986))
 - ▶ pNRQCD (weak coupling): $m_Q v \gg m_Q v^2, \Lambda_{QCD}$ (A. Pineda, JS, Nucl.Phys.Proc.Suppl.64:428-432,1998)
 - ▶ pNRQCD (strong coupling): $m_Q v, \Lambda_{QCD} \gg m_Q v^2$ (N. Brambilla, A. Pineda, JS, A. Vairo, Nucl.Phys.B566:275,2000)

How does the hadron look like ?



$$m_{QV} \sim 1/r \gg m_{QV}^2 \gtrsim \Lambda_{QCD}$$

weak coupling pNRQCD



$$m_{QV} \sim 1/r \gtrsim \Lambda_{QCD} \gg m_{QV}^2$$

strong coupling pNRQCD

|||
Born-Oppenheimer EFT

Figures: Najjar, Bali, 2009

pNRQCD weak coupling regime $\Lambda_{QCD} \lesssim m_Q v^2 \ll m_Q v$

$$\begin{aligned} \mathcal{L}_{\text{pNRQCD}} = \int d^3\mathbf{r} \text{Tr} \left\{ S^\dagger (i\partial_0 - h_s(\mathbf{r}, \mathbf{p}, \mathbf{P}_R, \mathbf{S}_1, \mathbf{S}_2, \mu)) S + \right. \\ \left. + O^\dagger (iD_0 - h_o(\mathbf{r}, \mathbf{p}, \mathbf{P}_R, \mathbf{S}_1, \mathbf{S}_2, \mu)) O \right\} \\ + V_A(r, \mu) \text{Tr} \{ O^\dagger \mathbf{r} \cdot g\mathbf{E} S + S^\dagger \mathbf{r} \cdot g\mathbf{E} O \} + \\ + \frac{V_B(r, \mu)}{2} \text{Tr} \{ O^\dagger \mathbf{r} \cdot g\mathbf{E} O + O^\dagger O \mathbf{r} \cdot g\mathbf{E} \} + \mathcal{O}(r^2, \frac{1}{m_Q}) \end{aligned}$$

- $h_{s,o} = \frac{\mathbf{p}^2}{m_Q} + V_{s,o}(r, \mu) + \mathcal{O}(\frac{1}{m_Q})$, quantum mechanical Hamiltonians with scale dependent potentials calculable in perturbation theory in $\alpha_s(m_Q v)$ and $1/m_Q$ ($V_s \simeq -4\alpha_s/3r$, $V_o \simeq \alpha_s/6r$)
- Spin symmetry holds in $h_{s,o}$ up to $\mathcal{O}(\frac{1}{m_Q^2})$
- $S=S(\mathbf{r}, \mathbf{R}, t)$, $O=O(\mathbf{r}, \mathbf{R}, t)$ are the color singlet/octet wave function fields
- $\mathbf{E}=\mathbf{E}(\mathbf{R}, t)$ is the chromoelectric field

Born-Oppenheimer EFT $m_Q v^2 \ll \Lambda_{\text{QCD}} \lesssim mv$

$$L_{\text{pNRQCD}} = \int d^3\mathbf{x}_1 \int d^3\mathbf{x}_2 S^\dagger (i\partial_0 - h_s(\mathbf{x}_1 - \mathbf{x}_2, \mathbf{p}_1, \mathbf{p}_2, \mathbf{S}_1, \mathbf{S}_2)) S,$$

$$h_s(\mathbf{x}_1, \mathbf{x}_2, \mathbf{p}_1, \mathbf{p}_2, \mathbf{S}_1, \mathbf{S}_2) = \frac{\mathbf{p}_1^2}{2m_Q} + \frac{\mathbf{p}_2^2}{2m_Q} + V_s(\mathbf{x}_1, \mathbf{x}_2, \mathbf{p}_1, \mathbf{p}_2, \mathbf{S}_1, \mathbf{S}_2),$$

$$V_s = V_s^{(0)} + \frac{V_s^{(1)}}{m_Q} + \frac{V_s^{(2)}}{m_Q^2} + \dots,$$

All V_s s can be, and most of them have been, calculated on the lattice

- $V_s^{(0)}$ and $V_s^{(1)}$ are central **Spin Symmetry holds**
- $V_s^{(2)}$ contains spin and velocity dependent terms

Born-Oppenheimer EFT at LO

- Matching to NRQCD in the static limit $\Rightarrow V_s^{(0)}$ is the ground state energy of two static color sources separated at a distance r
- Can be extracted from lattice calculations of the Wilson loop




Figure: Meyer, Swanson, 2015


- Well fitted by the Cornell potential

$$V_s^{(0)} = V_{\Sigma_g^+}(r) \approx -\frac{k_g}{r} + \kappa r + E_g^{Q\bar{Q}} \quad , \quad k_g = 0.489 \quad , \quad \kappa = 0.187 \text{ GeV}^2$$

BOEFT beyond LO

Ex. at $\mathcal{O}(1/m_Q^2)$: $V_{L_2 S_1}^{(1,1)}$ spin-orbit potential (Eichten, Feinberg, 79)

$$V_{L_2 S_1}^{(1,1)}(r) = -\frac{C_F}{r^2} i\mathbf{r} \cdot \lim_{T \rightarrow \infty} \int_0^T dt t \langle\langle g\mathbf{B}(t, \mathbf{r}/2) \times g\mathbf{E}(0, -\mathbf{r}/2) \rangle\rangle$$



$$V_2 = V_{L_2 S_1}^{(1,1)} / C_F, \text{ Koma, Koma, 09}$$

pNRQCD strong coupling regime beyond LO

An example at $\mathcal{O}(1/m_Q^2)$: the $V_{L_2 S_1}^{(1,1)}$ spin-orbit potential

- Short distance constraint: it must coincide with the perturbative evaluation,

$$V_{L_2 S_1}^{(1,1)}(r) \sim c_F \frac{C_F \alpha_s}{r^3}, \quad r \rightarrow 0$$

(Gupta, Radford, 81)

- Long distance constraint: it must coincide with the QCD effective string theory result

$$V_{L_2 S_1}^{(1,1)}(r) = -\frac{c_F g^2 \Lambda^2 \Lambda'}{\kappa r^2}, \quad r \rightarrow \infty$$

(Perez-Nadal, JS, 08)

$QQ/Q\bar{Q}$ + light quarks and gluons ($m_Q v, \Lambda_{QCD} \gg m_Q v^2$)

$$\mathcal{L}_{\text{HEH}} = \sum_{\kappa P} \Psi_{\kappa P}^\dagger [i\partial_t - h_{\kappa P}] \Psi_{\kappa P}$$

$$h_{\kappa P} = \frac{\mathbf{p}^2}{m_Q} + \frac{\mathbf{P}^2}{4m_Q} + V_{\kappa P}^{(0)}(\mathbf{r}) + \frac{1}{m_Q} V_{\kappa P}^{(1)}(\mathbf{r}, \mathbf{p}) + \mathcal{O}\left(\frac{1}{m_Q^2}\right)$$

- LDF \equiv light quarks + gluons, characterized by their quantum numbers ($\kappa, p \dots$)
 - ▶ $\kappa \equiv$ total angular momentum, $p \equiv$ parity (P)/CP
 - ▶ Quantum numbers not explicitly displayed: baryon number (B), isospin (I), strangeness (S), principal quantum number
- $V_{\kappa P}^{(0)}, V_{\kappa P}^{(1)}, \dots$ must be calculated non-perturbatively
- A truncation of \mathcal{L}_{HEH} needed for practical calculations \implies keep a limited number of lower lying κP

- $V_{\kappa P}^{(0)}$ is a $(2\kappa + 1) \times (2\kappa + 1) \times \mathbb{I}_2^{Q_1} \times \mathbb{I}_2^{Q_2}$ matrix, which can be decomposed into irreducible representations of $D_{\infty h}$, the symmetry group of a diatomic molecule

$$V_{\kappa P}^{(0)}(\mathbf{r}) = \sum_{\Lambda} V_{\kappa P \Lambda}^{(0)}(\mathbf{r}) \mathcal{P}_{\kappa \Lambda}$$

$\mathcal{P}_{\kappa \Lambda}$ projects onto LDF angular momenta $\pm \Lambda$ in the direction joining the two heavy quarks, $\Lambda = \kappa, \kappa - 1, \dots, \kappa - [\kappa]$

$$\mathcal{P}_{\frac{1}{2} \frac{1}{2}} = \mathbb{I}_2^{\text{lq}}$$

$$\mathcal{P}_{\frac{3}{2} \frac{1}{2}} = \frac{9}{8} \mathbb{I}_4^{\text{lq}} - \frac{1}{2} (\hat{\mathbf{r}} \cdot \mathbf{S}_{3/2})^2$$

$$\mathcal{P}_{\frac{3}{2} \frac{3}{2}} = -\frac{1}{8} \mathbb{I}_4^{\text{lq}} + \frac{1}{2} (\hat{\mathbf{r}} \cdot \mathbf{S}_{3/2})^2$$

$$\mathcal{P}_{10} = \mathbb{I}_3^{\text{lq}} - (\hat{\mathbf{r}} \cdot \mathbf{S}_1)^2$$

$$\mathcal{P}_{11} = (\hat{\mathbf{r}} \cdot \mathbf{S}_1)^2$$

...

- $V_{\kappa^P}^{(1)} = V_{\kappa^P\text{SI}}^{(1)} + V_{\kappa^P\text{SD}}^{(1)}$
- $V_{\kappa^P\text{SI}}^{(1)}$ does not depend on the spin or orbital angular momentum of the heavy quarks \implies admits the same decomposition as $V_{\kappa^P}^{(0)}$
- $V_{\kappa^P\text{SD}}^{(1)}$ depends on the spin and orbital angular momentum of the heavy quarks

$$V_{\kappa^P\text{SD}}^{(1)}(\mathbf{r}) = \sum_{\Lambda\Lambda'} \mathcal{P}_{\kappa\Lambda} \left[V_{\kappa^P\Lambda\Lambda'}^{sa}(\mathbf{r}) \mathbf{S}_{QQ} \cdot (\mathcal{P}_{10} \cdot \mathbf{S}_{\kappa}) + V_{\kappa^P\Lambda\Lambda'}^{sb}(\mathbf{r}) \mathbf{S}_{QQ} \cdot (\mathcal{P}_{11} \cdot \mathbf{S}_{\kappa}) \right. \\ \left. + V_{\kappa^P\Lambda\Lambda'}^l(\mathbf{r}) (\mathbf{L}_{QQ} \cdot \mathbf{S}_{\kappa}) \right] \mathcal{P}_{\kappa\Lambda'}$$

$$2\mathbf{S}_{QQ} = \boldsymbol{\sigma}_{QQ} = \boldsymbol{\sigma}_{Q_1} \mathbb{I}_{2Q_2} + \mathbb{I}_{2Q_1} \boldsymbol{\sigma}_{Q_2} \quad , \quad \mathcal{P}_{10}^{ij} = \hat{\mathbf{r}}^i \hat{\mathbf{r}}^j \quad , \quad \mathcal{P}_{11}^{ij} = \delta^{ij} - \hat{\mathbf{r}}^i \hat{\mathbf{r}}^j$$

Matching to NRQCD

- Build an NRQCD operator with the quantum numbers of $\Psi_{\kappa P}$

$$\mathcal{O}_{\kappa P}^{Q\bar{Q}}(t, \mathbf{r}, \mathbf{R}) = \chi_c^\top(t, \mathbf{x}_2) \phi(t, \mathbf{x}_2, \mathbf{R}) \mathcal{Q}_{Q\bar{Q}\kappa P}(t, \mathbf{R}) \phi(t, \mathbf{R}, \mathbf{x}_1) \psi(t, \mathbf{x}_1)$$

$$\mathcal{O}_{\kappa P}^{QQ}(t, \mathbf{r}, \mathbf{R}) = \psi^\top(t, \mathbf{x}_2) \phi^\top(t, \mathbf{R}, \mathbf{x}_2) \mathcal{Q}_{QQ\kappa P}(t, \mathbf{R}) \phi(t, \mathbf{R}, \mathbf{x}_1) \psi(t, \mathbf{x}_1)$$

- Examples:

- ▶ Hybrid

$$\mathcal{Q}_{1+-}^\alpha(t, \mathbf{x}) = (\mathbf{e}_\alpha^\dagger \cdot \mathbf{B}(t, \mathbf{x}))$$

- ▶ $Q\bar{Q}q\bar{q}$ tetraquark

$$\mathcal{Q}_{0++}(t, \mathbf{x}) = [\bar{q}(t, \mathbf{x}) T^a q(t, \mathbf{x})] T^a$$

- ▶ Doubly heavy baryons

$$\mathcal{Q}_{(1/2)+}^\alpha(t, \mathbf{x}) = \underline{T}^I [P_+ q^I(t, \mathbf{x})]^\alpha$$

- ▶ $QQ\bar{q}\bar{q}$ tetraquark

$$\mathcal{Q}_{0-}(t, \mathbf{x}) = [\bar{q}(t, \mathbf{x}) \underline{T}^I \gamma^2 q^*(t, \mathbf{x})] \underline{T}^I$$


Matching to NRQCD

- Impose $\mathcal{O}_{\kappa^P}^h(t, \mathbf{r}, \mathbf{R}) = \sqrt{Z_{h\kappa^P}} \Psi_{h\kappa^P}(t, \mathbf{r}, \mathbf{R}), \quad h = QQ, Q\bar{Q}.$

$$\langle 0 | T \{ \mathcal{O}_{\kappa^P}^h(t/2) \mathcal{O}_{\kappa^P}^{h\dagger}(-t/2) \} | 0 \rangle = \sqrt{Z_{h\kappa^P}} \langle 0 | T \{ \Psi_{h\kappa^P}(t/2) \Psi_{h\kappa^P}^\dagger(-t/2) \} | 0 \rangle \sqrt{Z_{h\kappa^P}^\dagger}$$

- ▶ Then at $\mathcal{O}(1)$

$$V_{h\kappa^P\Lambda}^{(0)}(\mathbf{r}) = \lim_{t \rightarrow \infty} \frac{i}{t} \log \left(\text{Tr} \left[\mathcal{P}_{\kappa\Lambda} \langle 1 \rangle_{\square}^{h\kappa^P} \right] \right)$$



- ▶ At $\mathcal{O}\left(\frac{1}{m_Q}\right)$, for instance,

$$\begin{aligned}
 V_{\kappa^p \Lambda \Lambda'}^{sb} &= -c_F \lim_{t \rightarrow \infty} \sqrt{\frac{\text{Tr}[\mathcal{P}_{\kappa \Lambda}] \text{Tr}[\mathcal{P}_{\kappa \Lambda'}]}{\text{Tr}[\mathcal{P}_{\kappa \Lambda} \langle 1 \rangle_{\square}^{h\kappa^p}] \text{Tr}[\mathcal{P}_{\kappa \Lambda'} \langle 1 \rangle_{\square}^{h\kappa^p}]}} \\
 &\times \frac{\ln\left(\frac{\text{Tr}[\mathcal{P}_{\kappa \Lambda} \langle 1 \rangle_{\square}^{h\kappa^p}] \text{Tr}[\mathcal{P}_{\kappa \Lambda'}]}{\text{Tr}[\mathcal{P}_{\kappa \Lambda'} \langle 1 \rangle_{\square}^{h\kappa^p}] \text{Tr}[\mathcal{P}_{\kappa \Lambda}]}\right)}{2t \sinh\left(\ln \sqrt{\frac{\text{Tr}[\mathcal{P}_{\kappa \Lambda} \langle 1 \rangle_{\square}^{h\kappa^p}] \text{Tr}[\mathcal{P}_{\kappa \Lambda'}]}{\text{Tr}[\mathcal{P}_{\kappa \Lambda'} \langle 1 \rangle_{\square}^{h\kappa^p}] \text{Tr}[\mathcal{P}_{\kappa \Lambda}]}}\right)} \\
 &\times \int_{-t/2}^{t/2} dt' \frac{\text{Tr}\left[\left(\mathbf{S}_{\kappa} \cdot \mathcal{P}_{11}\right) \cdot \left(\mathcal{P}_{\kappa \Lambda} \langle \mathbf{g} \mathbf{B}(t', \mathbf{x}_1) \rangle_{\square}^{h\kappa^p} \mathcal{P}_{\kappa \Lambda'}\right)\right]}{\text{Tr}\left[\left(\mathbf{S}_{\kappa} \cdot \mathcal{P}_{11}^{\text{c.r.}}\right) \cdot \left(\mathcal{P}_{\kappa \Lambda} \mathbf{S}_{\kappa} \mathcal{P}_{\kappa \Lambda'}\right)\right]}
 \end{aligned}$$

Applications

- Doubly Heavy Baryons: QQq (JS, Tarrús Castellà, 20, 21; see my talk on Thu)
- Hyperfine splittings of Heavy Quarkonium Hybrids: $Q\bar{Q}g$ (JS, Tomàs Valls, 23)

Disclaimer:

- Interactions with heavy-light meson/baryon pairs neglected
- They have been recently addressed in the BOEFT for heavy quarkonium (Tarrús Castellà, 22)
- It can be easily generalized to heavy exotics

Heavy Quarkonium Hybrids

- See Abhishek's talk last Wed
- Spin average spectrum (BO approximation; Braaten, Langmack, Hudson Smith, 2014; Berwin, Brambilla, Tarrús Castellà, Vairo, 15; Oncala, Soto, 17)
 - ▶ Based on lattice data (Juge, Kuti, Morningstar, 02; Bali, Pineda, 03)
 - ▶ More recent and accurate lattice data available (Capitani, Philipsen, Reisinger, Riehl, Wagner, 18; Schlosser, Wagner, 21)
- Inclusive decay width to heavy quarkonium (Oncala, JS, 17)
 - ▶ Revisited in (Brambilla, Lai, Mohapatra, Vairo, 22)
 - ★ Improved the $\Delta S = 0$ transitions $\mathcal{O}(1/m_Q^0)$
 - ★ Calculated the $\Delta S = 1$ transitions $\mathcal{O}(1/m_Q^2)$
- Mixing with heavy quarkonium ($\mathcal{O}(1/m_Q)$, spin-dependent; Oncala, JS, 17)
 - ▶ Important effects when a quarkonium state and a hybrid state with the same quantum numbers have similar masses
 - ▶ Leads to violations of spin conservation
 - ▶ Likely to increase the estimates of the $\Delta S = 1$ transitions above

The hyperfine splitting of heavy quarkonium hybrids

- The lower lying hybrid potentials correspond to $\kappa^P = 1^+$
- This leads to two spin projections on the direction $Q-\bar{Q}$, $\Lambda = 0, 1$
- The general formulas above imply that there are two independent potentials: $V_{1^+11}^{sa}(r)$, $V_{1^+10}^{sb}(r)$ (Oncala, JS, 17; Brambilla, Lai, Segovia, Tarrús Castellà, Vairo, 18, 19)
- No lattice calculation available for them. How to estimate them?
 - ▶ Brambilla et al. used weak coupling pNRQCD short distance expressions to estimate them which hold for $r \ll 1/\Lambda_{QCD}$. The $1/m_Q^2$ spin dependent potentials were also included.
 - ▶ We (JS, Tomàs Valls, 23) use an interpolation between the short distance expressions and long distance ones calculated in the QCD effective string theory (Pérez-Nadal, JS, 08; Brambilla, Groher, Martinez, Vairo, 14).
- Typical values: $\langle 1/r \rangle \sim 0.17 - 0.42$ GeV for $c\bar{c}g$, $\langle 1/r \rangle \sim 0.22 - 0.53$ GeV for $b\bar{b}g$ (Berwein, Brambilla, Tarrús Castellà, Vairo, 14)

Hyperfine Splittings

JS, 17

- They appear at $\mathcal{O}(1/m_Q)$ ($\mathcal{O}(1/m_Q^2)$) in hybrids (quarkonium)
- They lead to the following mass formulae

$$\frac{M_{1J+1} - M_{0J}}{M_{1J} - M_{0J}} = -J \quad \frac{M_{1J-1} - M_{0J}}{M_{1J} - M_{0J}} = J + 1$$

$$(s/d)_1 : M_{2-+} + M_{0-+} = M_{1-+} + M_{1--}$$

$$p_1 : M_{2+-} + M_{0+-} = M_{1+-} + M_{1++}$$

$$(p/f)_2 : M_{3+-} + M_{1+-} = M_{2+-} + M_{2++}$$

$$d_2 : M_{3-+} + M_{1-+} = M_{2-+} + M_{2--}$$

- Consistent with the values of the lattice **HSC**
- Induces mixing between different hybrid states

The short distance potentials

- The two independent potentials are rearranged in

$$V_{hf}(r) = \frac{1}{6} V_{1+11}^{sa}(r) - \frac{1}{3} V_{1+10}^{sa}(r) \quad (\text{spin} - \text{spin})$$

$$V_{hf2}(r) = -\frac{1}{2} \left(V_{1+11}^{sa}(r) + V_{1+10}^{sb}(r) \right) \quad (\text{tensor})$$

- At short distances:

$$\begin{aligned} V_{hf}(r)/m_Q &= A + \mathcal{O}(r^2) & , & \quad A \sim c_F \Lambda_{QCD}^2 / m_Q \\ V_{hf2}(r)/m_Q &= B r^2 + \mathcal{O}(r^4) & , & \quad B \sim c_F \Lambda_{QCD}^4 / m_Q \end{aligned}$$

The short distance potentials depend on two unknown non-perturbative parameters.

The long distance potentials

- At long distances:

$$\frac{V_{1+11}^{sa}(r)}{m_Q} = -\frac{2c_F\pi^2 g\Lambda'''}{m_Q\kappa r^3} \equiv V_{ld}^{sa}(r)$$

$$\frac{V_{1+10}^{sb}(r)}{m_Q} = \mp \frac{c_F g\Lambda'\pi^2}{m_Q\sqrt{\pi\kappa}} \frac{1}{r^2} \equiv V_{ld}^{sb}(r)$$

- ▶ κ is the string tension $\sim \Lambda_{QCD}^2$
- ▶ $g\Lambda'$, $g\Lambda''' \sim \Lambda_{QCD}$ also enter the spin dependent potentials for heavy quarkonium
- ▶ They can be extracted from lattice calculations of those potentials (Koma, Koma, 09)

$$g\Lambda' \sim -59 \text{ MeV} \quad ; \quad g\Lambda''' \sim \pm 230 \text{ MeV}$$

(Oncala, JS, 17)

The interpolating potentials

- We use the following interpolation

$$\frac{V_{hf}(r)}{m_Q} = \frac{A + \left(\frac{r}{r_0}\right)^2 \left(\frac{1}{6} V_{ld}^{sa}(r_0) - \frac{r}{3r_0} V_{ld}^{sb}(r_0)\right)}{1 + \left(\frac{r}{r_0}\right)^5}$$

$$\frac{V_{hf2}(r)}{m_Q} = \frac{Br^2 - \left(\frac{r}{r_0}\right)^5 \left(\frac{r_0}{2r} V_{ld}^{sa}(r_0) + \frac{1}{2} V_{ld}^{sb}(r_0)\right)}{1 + \left(\frac{r}{r_0}\right)^7}.$$

- $r_0 \simeq 3.96 \text{ GeV}^{-1} \sim 1/\Lambda_{QCD}$

Charmonium Hybrids HFS

- We use lattice data of the HSC for charmonium to fix A and B (relativistic charm, $m_\pi \sim 240$ MeV, Cheung, O'Hara, Moir, Peardon, Ryan, Thomas, Tims, 16)
- We focus on hyperfine splittings not on spin averages
- Same strategy as Brambilla et al. , 19
 - ▶ We have a 2 parameter fit and get $A = 0.115 \pm 0.034$ GeV, $B = 0.0038 \pm 0.0154$ GeV³ with a $\chi^2/\text{dof} \sim 0.64$
 - ▶ Brambilla et al. , 19 have an 8 parameter fit with a $\chi^2/\text{dof} \sim 0.99$
- Including long distance information from the QCD string improves the description of lattice data

Charmonium Hybrids HFS




Figure: The spectrum of the lower-lying $n(s/d)_1$ (H_1), np_1 (H_2), $n(p/f)_2$ (H_4) and np_0 (H_3) charmonium hybrids

Bottomonium Hybrids HFS




Figure: The spectrum of the lower-lying $n(s/d)_1$ (H_1), np_1 (H_2), $n(p/f)_2$ (H_4) and np_0 (H_3) bottomonium hybrids

Conclusions

- The BOEFT provides a QCD based framework to address doubly-heavy exotics systematically
- It requires non-perturbative potentials as an input
- When those potentials are not available, a Cornell-like approach of interpolating between the short distance QCD (pNRQCD) calculation and a long distance string calculation appears to be promising